

1 Approximating activation edge-cover and facility 2 location problems

3 Zeev Nutov

4 The Open University of Israel

5 nutov@openu.ac.il

6 Guy Kortsarz

7 Rutgers University, Camden

8 guyk@camden.rutgers.edu

9 Eli Shalom¹

10 The Open University of Israel

11 eli.shalom@gmail.com

12 — Abstract —

13 What approximation ratio can we achieve for the FACILITY LOCATION problem if whenever a client
14 u connects to a facility v , the opening cost of v is at most θ times the service cost of u ? We
15 show that this and many other problems are a particular case of the ACTIVATION EDGE-COVER
16 problem. Here we are given a multigraph $G = (V, E)$, a set $R \subseteq V$ of terminals, and thresholds
17 $\{t_u^e, t_v^e\}$ for each uv -edge $e \in E$. The goal is to find an assignment $\mathbf{a} = \{a_v : v \in V\}$ to the nodes
18 minimizing $\sum_{v \in V} a_v$, such that the edge set $E_{\mathbf{a}} = \{e = uv : a_u \geq t_u^e, a_v \geq t_v^e\}$ activated by \mathbf{a}
19 covers R . We obtain ratio $1 + \omega(\theta) \approx \ln \theta - \ln \ln \theta$ for the problem, where $\omega(\theta)$ is the root of the
20 equation $x + 1 = \ln(\theta/x)$ and θ is a problem parameter. This result is based on a simple generic
21 algorithm for the problem of minimizing a sum of a decreasing and a sub-additive set functions,
22 which is of independent interest. As an application, we get that the above variant of FACILITY
23 LOCATION admits ratio $1 + \omega(\theta)$; if for each facility all service costs are identical then we show a
24 better ratio $1 + \max_{k \geq 1} \frac{H_k - 1}{1 + k/\theta}$, where $H_k = \sum_{i=1}^k 1/i$. For the MIN-POWER EDGE-COVER problem
25 we improve the ratio 1.406 of [3] (achieved by iterative randomized rounding) to $1 + \omega(1) < 1.2785$.
26 For unit thresholds we improve the ratio $73/60 \approx 1.217$ of [3] to $\frac{1555}{1347} \approx 1.155$.

27 **2012 ACM Subject Classification** Mathematics of computing \rightarrow Approximation algorithms

28 **Keywords and phrases** generalized min-covering problem, activation edge-cover, facility location,
29 minimum power, approximation algorithm

30 **Digital Object Identifier** 10.4230/LIPIcs.CVIT.2016.23

31 **Funding** *Guy Kortsarz*: This work was partially supported by NSF grant number 434923.

32 **1** Introduction

33 Let $G = (V, E)$ be an undirected multigraph where each edge $e \in E$ has an **activating**
34 **function** f^e from some range $L^e \subseteq \mathbb{R}_+^2$ to $\{0, 1\}$. Given a non-negative **assignment**
35 $\mathbf{a} = \{a_v : v \in V\}$ to the nodes, we say that a uv -edge $e \in E$ is **activated** by \mathbf{a} if
36 $f^e(a_u, a_v) = 1$. Let $E_{\mathbf{a}} = \{e \in E : f^e(a_u, a_v) = 1\}$ denote the set of edges activated by
37 \mathbf{a} . The **value** of an assignment \mathbf{a} is $\mathbf{a}(V) = \sum_{v \in V} a_v$. In ACTIVATION NETWORK DESIGN
38 problems the goal is to find an assignment \mathbf{a} of minimum value, such that the edge set $E_{\mathbf{a}}$
39 activated by \mathbf{a} satisfies a prescribed property. We refer the reader to a paper of Panigrahi [16]
40 and a recent survey [15] on activation problems, where also the following two assumptions
41 are justified.

¹ Part of this work was done as a part of author's M.Sc. thesis done at the Open University of Israel.



23:2 Approximating activation edge-cover and facility location problems

42 **Monotonicity Assumption.** For every $e \in E$, f^e is monotone non-decreasing, namely,
 43 $f^e(x_u, x_v) = 1$ implies $f^e(y_u, y_v) = 1$ if $y_u \geq x_u$ and $y_v \geq x_v$.

44 **Polynomial Domain Assumption.** Every $v \in V$ has a polynomial size in $n = |V|$ set L_v
 45 of “levels” and $L^e = L_u \times L_v$ for every uv -edge $e \in E$.

46 Given a set $R \subseteq V$ of **terminals** we say that an edge set J is an R -**cover** or that J
 47 **covers** R if every $v \in R$ has some edge in J incident to it. In the EDGE-COVER problem
 48 we seek an R -cover of minimum value. The min-cost version of this problem can be solved
 49 in polynomial time [7], and it is one of the most fundamental problems in Combinatorial
 50 Optimization, cf. [19].

51 We consider the ACTIVATION EDGE-COVER problem. Since we consider multigraphs,
 52 $e = uv$ means that e is a uv -edge, namely, that u, v are the endnodes of e ; $e = uv \in E$ means
 53 that $e \in E$ is a uv -edge. Under the two assumptions above, the problem can be can be
 54 formulated without activating functions. For this, replace each edge $e = uv$ by a set of at
 55 most $|L_u| \cdot |L_v|$ uv -edges $\{e(t_u, t_v) : (t_u, t_v) \in L_u \times L_v, f^e(t_u, t_v) = 1\}$. Then for any $J \subseteq E$
 56 the optimal assignment \mathbf{a} activating J is given by $a_u = \max\{t_u^e : e \in J \text{ is incident to } u\}$;
 57 here and everywhere a maximum or a minimum taken over an empty set is assumed to be
 58 zero. Consequently, the problem can be restated as follows.

ACTIVATION EDGE-COVER

59 **Input:** A graph $G = (V, E)$, a set of terminals $R \subseteq V$, and thresholds $\{t_u^e, t_v^e\}$ for each
 uv -edge $e \in E$.

Output: An assignment \mathbf{a} of minimum value $\mathbf{a}(V) = \sum_{v \in V} a_v$, such that the edge set
 $E_{\mathbf{a}} = \{e = uv \in E : a_u \geq t_u^e, a_v \geq t_v^e\}$ activated by \mathbf{a} covers R .

60 As we will explain later, ACTIVATION EDGE-COVER problems are among the most
 61 fundamental problems in network design, that include NP-hard problems such as SET-
 62 COVER, FACILITY LOCATION, covering problems that arise in wireless networks (node
 63 weighted/min-power/installation problems), and many other problems.

64 To state our main result we define assignments \mathbf{q} and \mathbf{c} , where $c_v = q_v = 0$ if $v \in V \setminus R$
 65 and for $u \in R$:

- 66 • $q_u = \min_{e=uv \in E} t_u^e$ is the minimum threshold at u of an edge in E incident to u .
- 67 • $c_u = \min_{e=uv \in E} (t_u^e + t_v^e) - q_u$, so $c_u + q_u$ is the minimum value of an edge in E incident to u .

The quantity $\max_{u \in R} c_u/q_u$ is called the **slope** of the instance. We say that an ACTIVATION
 EDGE-COVER instance is θ -**bounded** if the instance slope is at most θ , namely if $c_u \leq \theta q_u$
 for all $u \in R$; moreover, we assume by default that $\theta = \max_{u \in R} c_u/q_u$ is the instance slope.
 For each $u \in R$ let e_u be some minimum value edge covering u . Then $\{e_u : u \in R\}$ is an
 R -cover of value at most $(\mathbf{c} + \mathbf{q})(R)$. From this and the definition of θ we get

$$0 \leq \text{opt} - \mathbf{q}(R) \leq \mathbf{c}(R) \leq \theta \mathbf{q}(R) \leq \theta \text{opt}$$

68 In particular, $(\mathbf{c} + \mathbf{q})(R) \leq (\theta + 1)\text{opt}$. Using this, it is possible to design a greedy algorithm
 69 with ratio $1 + \ln(\theta + 1)$. We will show how to obtain a better ratio (the difference is quite
 70 significant when $\theta \leq 10^4$ – see Table 1), as follows.

71 The **Lambert W -Function** (a.k.a. ProductLog Function) $W(z)$ is the inverse function
 72 of $f(W) = We^W$. It is known that for any $\theta > 0$, $\omega(\theta) = W(\theta/e)$ equals to the (unique) real
 73 root of the equation $x + 1 = \ln(\theta/x)$, and that $\lim_{\theta \rightarrow \infty} [1 + \omega(\theta) - (\ln \theta - \ln \ln \theta)] = 0$. Our
 74 main result is:

θ	1	2	3	4	5	10	100	1000	10000	1000000
$1 + \omega(\theta)$	1.2785	1.4631	1.6036	1.7179	1.8146	2.1569	3.6360	5.4214	7.3603	11.4673
$1 + \bar{\omega}(\theta)$	1.2167	1.3667	1.4834	1.5800	1.6637	1.9645	3.3428	5.0808	6.9967	11.0820
$\ln \theta - \ln \ln \theta$	-	1.0597	1.0046	1.0597	1.1336	1.4686	3.0780	4.9752	6.9901	11.1898
$1 + \ln(\theta + 1)$	1.6932	2.0987	2.3863	2.6095	2.7918	3.3979	5.6152	7.9088	10.2105	14.8156

■ **Table 1** Some numerical bounds on $1 + \omega(\theta)$, $1 + \bar{\omega}(\theta)$, $\ln \theta - \ln \ln \theta$, and $1 + \ln(\theta + 1)$.

75 ▶ **Theorem 1.** ACTIVATION EDGE-COVER admits ratio $1 + \omega(\theta)$ for θ -bounded instances.
 76 The problem also admits ratio $1 + \ln(\Delta + 1)$, and ratio $1 + \ln \Delta$ if R is an independent set in
 77 G , where Δ is the maximum number of terminal neighbors of a node in G .

78 This result is based on a generic simple approximation algorithm for the problem of
 79 minimizing a sum of a decreasing and a sub-additive set functions, which is of independent
 80 interest; it is described in the next section. This result is inspired by the so called “Loss
 81 Contraction Algorithm” of Robins & Zelikovsky [18] for the STEINER TREE problem, and
 82 the analysis in [10] of this algorithm.

83 Let us say that $v \in V$ is a **steady node** if the thresholds t_e^v of the edges e incident to
 84 v are all equal the same number w_v , which we call the **weight of v** . Note that we may
 85 assume that all non-terminals are steady, by replacing each $v \in V \setminus R$ by L_v new nodes; see
 86 the so called “Levels Reduction” in [15]. This implies that we may also assume that no two
 87 parallel edges are incident to the same non-terminal. Clearly, we may assume that $R \setminus V$ is
 88 an independent set in G . Let BIPARTITE ACTIVATION EDGE-COVER be the restriction of
 89 ACTIVATION EDGE-COVER to instances when also R is an independent set, namely, when G is
 90 bipartite with sides $R, V \setminus R$. Note that in this case G is a simple graph and all non-terminals
 91 are steady.

92 We now mention some particular threshold types in ACTIVATION EDGE-COVER problems,
 93 some known problems arising from these types, and some implications of Theorem 1 for these
 94 problems.

95 WEIGHTED SET-COVER

96 This is a particular case of BIPARTITE ACTIVATION EDGE-COVER when all nodes are steady
 97 and nodes in R have weight 0. Note that in this case θ is infinite, and we can only deduce
 98 from Theorem 1 the known ratio $1 + \ln \Delta$. Consider a modification of the problem, which we
 99 call θ -BOUNDED WEIGHTED SET-COVER: when we pick a set $v \in V \setminus R$, we need to pay
 100 w_v/θ for each element in R covered by v . Then the corresponding ACTIVATION EDGE-COVER
 101 instance is θ -bounded.

102 FACILITY LOCATION

103 Here we are given a bipartite graph with sides R (clients) and $V \setminus R$ (facilities), weights
 104 (opening costs) $\mathbf{w} = \{w_v : v \in V \setminus R\}$, and distances (service costs) $\mathbf{d} = \{d_{uv} : u \in$
 105 $R, v \in V \setminus R\}$. We need to choose $S \subseteq V \setminus R$ with $\mathbf{w}(S) + \sum_{u \in R} d(u, S)$ minimal, where
 106 $d(u, S) = \min_{v \in S} d_{uv}$ is the minimal distance from u to S . This is equivalent to BIPARTITE
 107 ACTIVATION EDGE-COVER. Note however that if for some constant θ we have $w_v \leq \theta d_{uv}$
 108 for all $uv \in E$ with $u \in R$ and $v \in V \setminus R$, then the corresponding BIPARTITE ACTIVATION
 109 EDGE-COVER instance is θ -bounded, and achieves a low constant ratio even for large values
 110 of θ .

111 INSTALLATION EDGE-COVER

112 Suppose that the installation cost of a wireless network is proportional to the total height of
 113 the towers for mounting antennas. An edge uv is activated if the towers at u and v are tall

23:4 Approximating activation edge-cover and facility location problems

114 enough to overcome obstructions and establish line of sight between the antennas. This is
 115 modeled as each pair $u, v \in V$ has a height demand h^{uv} and constants γ_{uv}, γ_{vu} , such that
 116 a uv -edge is activated by \mathbf{a} if the scaled heights $\gamma_{uv}a_u, \gamma_{vu}a_v$ sum to at least h^{uv} . In the
 117 INSTALLATION EDGE-COVER problem, we need to assign heights to the antennas such that
 118 each terminal can communicate with some other node, while minimizing the total sum of the
 119 heights. The problem is SET-COVER hard even for 0, 1 thresholds and bipartite G [16]. But
 120 in a practical scenario, the quotient of the maximum tower height over the minimum tower
 121 height is usually bounded by a constant; say, if possible tower heights are 5, 15, 20, then the
 122 slope is $\theta = 4$.

123 MIN-POWER EDGE-COVER

124 This problem is a particular case of ACTIVATION EDGE-COVER when $t_u^e = t_v^e$ for every edge
 125 $e = uv \in E$; note that $\theta = 1$ in this case (in fact, the case $\theta = 1$ is much more general). The
 126 motivation is to assign energy levels to the nodes of a wireless network while minimizing the
 127 total energy consumption, and enabling communication for every terminal. The MIN-POWER
 128 EDGE-COVER problem is NP-hard even if $R = V$, or if R is an independent set in the input
 129 graph G and unit thresholds [11]. The problem admits ratio 2 by a trivial reduction to the
 130 min-cost case. This was improved to 1.5 in [13], and then to 1.406 in [3], where is also given
 131 ratio $73/60$ for the bipartite case and for unit thresholds.

132 From Theorem 1 and the discussion above we get:

133 ► **Corollary 2.** MIN-POWER EDGE-COVER admits ratio $1 + \omega(1) < 1.2785$, and the θ -bounded
 134 versions of each one of the problems WEIGHTED SET-COVER, FACILITY LOCATION, and
 135 INSTALLATION EDGE-COVER, admits ratio $1 + \omega(\theta)$.

136 Let us illustrate this result on the FACILITY LOCATION problem. One might expect a
 137 constant ratio for any $\theta > 0$, but our ratio $1 + \omega(\theta)$ is surprisingly low. Even if $\theta = 100$
 138 (service costs are at least 1% of opening costs) then we get a small ratio $1 + \omega(100) < 3.636$.
 139 Even for $\theta = 10^4$ we still get a reasonable ratio $1 + \omega(10^4) < 7.3603$. All previous results for
 140 the problem are usually summarized by just two observations: the problem is SET-COVER
 141 hard (so has a logarithmic approximation threshold by [17, 8]), and that it admits a matching
 142 logarithmic ratio $1 + \ln |R|$ [5]; see surveys on FACILITY LOCATION problems by Vygen [22]
 143 and Shmoys [20]. Due to this, all work focused on the more tractable METRIC FACILITY
 144 LOCATION problem. Our Theorem 1 implies that many practical non-metric FACILITY
 145 LOCATION instances admit a reasonable small constant ratio.

146 For the case of “locally uniform” thresholds – when for each non-terminal (facility) all
 147 thresholds (service costs) are identical, we show a better ratio, see also Table 1. In what
 148 follows, let H_k denote the k -th harmonic number.

149 ► **Theorem 3.** BIPARTITE ACTIVATION EDGE-COVER with locally uniform thresholds admits
 150 ratio $1 + \bar{\omega}(\theta)$, where $\bar{\omega}(\theta) = \max_{k \geq 1} \frac{H_k - 1}{1 + k/\theta}$.

151 We do not have a convenient formula for $\bar{\omega}(\theta)$, but in Section 4 we observe that the
 152 maximum is attained for the smallest integer k_θ such that $H_{k_\theta} \geq 2 + \frac{\theta-1}{k_\theta+1}$. We will show
 153 that $\bar{\omega}(\theta) < \omega(\theta)$ for all θ , and that $\lim_{\theta \rightarrow \infty} [\omega(\theta) - \bar{\omega}(\theta)] = 0$, so both $\omega(\theta)$ and $\bar{\omega}(\theta)$ are
 154 close to $\ln \theta - \ln \ln \theta$ for large values of θ , although the convergence is very slow; see also
 155 Table 1. We note that Slavik [21] proved that the greedy algorithm for SET-COVER achieves
 156 ratio $\ln n - \ln \ln n + \Theta(1)$, while our ratio for ACTIVATION EDGE-COVER is asymptotically
 157 $\ln \theta - \ln \ln \theta + \Theta(1)$; but we do not see an immediate relation between the two results.

158 We will also show that $\bar{\omega}(1) = 73/60$. Note that our Theorem 1 ratio 1.2785 for $\theta = 1$
 159 significantly improves the previous best ratio 1.406 of [3] for MIN-POWER EDGE-COVER on
 160 general graphs achieved by iterative randomized rounding; we do not match the ratio 73/60
 161 of [3] for the bipartite case, but note that the case $\theta = 1$ is much more general than the
 162 min-power case considered in [3].

163 In addition, we consider unit thresholds, and using some ideas from [3] improve the
 164 previous best ratio 73/60 of [3] as follows.

165 ► **Theorem 4.** ACTIVATION EDGE-COVER with unit thresholds admits ratio $\frac{1555}{1347}$.

166 We note that our main contribution is not technical, although some proofs are non-
 167 trivial (the reader may observe that proofs of many seemingly complicated results were
 168 substantially simplified with years, by additional effort). Our main contribution is giving a
 169 unified algorithm for a large class of problems that we identify – θ -BOUNDED ACTIVATION
 170 EDGE-COVER problems, either substantially improving known ratios, or showing that many
 171 seemingly SET-COVER hard problems may be tractable in practice. Let us also point out that
 172 our main result is more general than the applications listed in Corollary 2. The generalization
 173 to θ -bounded ACTIVATION EDGE-COVER problems is different from earlier results; besides
 174 finding a unifying algorithmic idea generalizing and improving previous results, we are also
 175 able to find tractable special cases in a new direction.

176 The rest of this paper is organized as follows. In Section 2 we define the GENERALIZED
 177 MIN-COVERING problem and analyze a greedy algorithm for it, see Theorem 5. In Section 3
 178 we use Theorem 5 to prove Theorem 1. Theorems 3 and 4 are proved using a modified
 179 method in Sections 4 and 5, respectively.

180 2 The GENERALIZED MIN-COVERING problem

181 A set function f is **increasing** if $f(A) \leq f(B)$ whenever $A \subseteq B$; f is **decreasing** if $-f$ is
 182 increasing, and f is **sub-additive** if $f(A \cup B) \leq f(A) + f(B)$ for any subsets A, B of the
 183 ground-set. Let us consider the following algorithmic problem:

GENERALIZED MIN-COVERING

184 *Input:* Non-negative set functions ν, τ on subsets of a ground-set U such that ν is
 decreasing, τ is sub-additive, and $\tau(\emptyset) = 0$.

Output: $A \subseteq U$ such that $\nu(A) + \tau(A)$ is minimal.

185 The “ordinary” MIN-COVERING problem is $\min\{\tau(A) : \nu(A) = 0\}$; it is a particular case
 186 of the GENERALIZED MIN-COVERING problem when we seek to minimize $M\nu(A) + \tau(A)$ for
 187 a large enough constant M . Under certain assumptions, the MIN-COVERING problem admits
 188 ratio $1 + \ln \nu(\emptyset)$ [12]. Various generic covering problems are considered in the literature,
 189 among them the SUBMODULAR COVERING problem [23], and several other types, cf. [4].
 190 The variant we consider is inspired by the algorithms of Robins & Zelikovsky [18] for the
 191 STEINER TREE problem, and the analysis in [10] of this algorithm; but, to the best of our
 192 knowledge, the explicit formulation of the GENERALIZED MIN-COVERING problem given
 193 here is new. Interestingly, our ratio for MIN-POWER EDGE-COVER is the same as that of
 194 [18] for STEINER TREE in quasi-bipartite graphs.

195 We call ν the **potential** and τ the **payment**. The idea behind this interpretation and
 196 the subsequent greedy algorithm is as follows. Given an optimization problem, the potential
 197 $\nu(A)$ is the value of some “simple” augmenting feasible solution for A . We start with an

198 empty set solution, and iteratively try to decrease the potential by adding a set $B \subseteq U \setminus A$
 199 of minimum “density” – the price paid for a unit of the potential. The algorithm terminates
 200 when the price ≥ 1 , since then we gain nothing from adding B to A . The ratio of such an
 201 algorithm is bounded by $1 + \ln \frac{\nu(\emptyset)}{\text{opt}}$ (assuming that during each iteration a minimum density
 202 set can be found in polynomial time). So essentially the greedy algorithm converts ratio
 203 $\alpha = \frac{\nu(\emptyset)}{\text{opt}}$ into ratio $1 + \ln \alpha$. However, sometimes a tricky definition of the potential and the
 204 payment functions may lead to a smaller ratio.

205 Let opt be the optimal solution value of a problem instance at hand. Fix an optimal
 206 solution A^* . Let $\nu^* = \nu(A^*)$, $\tau^* = \tau(A^*)$, so $\text{opt} = \tau^* + \nu^*$. The quantity $\frac{\tau(B)}{\nu(A) - \nu(A \cup B)}$ is
 207 called the **density** of B (w.r.t. A); this is the price paid by B for a unit of potential. The
 208 GREEDY ALGORITHM (a.k.a. Relative Greedy Heuristic) for the problem starts with $A = \emptyset$
 209 and while $\nu(A) > \nu^*$ repeatedly adds to A a non-empty augmenting set B that satisfies the
 210 following condition, while such B exists:

$$211 \quad \text{Density Condition:} \quad \frac{\tau(B)}{\nu(A) - \nu(A \cup B)} \leq \min \left\{ 1, \frac{\tau^*}{\nu(A) - \nu^*} \right\}.$$

212 Note that since ν is decreasing $\nu(A) - \nu(A \cup A^*) \geq \nu(A) - \nu(A^*) = \nu(A) - \nu^*$; hence if
 213 $\nu(A) > \nu^*$, then $\frac{\tau(A^*)}{\nu(A) - \nu(A \cup A^*)} \leq \frac{\tau^*}{\nu(A) - \nu^*}$ and there exists an augmenting set B that satisfies
 214 the condition $\frac{\tau(B)}{\nu(A) - \nu(A \cup B)} \leq \frac{\tau^*}{\nu(A) - \nu^*}$, e.g., $B = A^*$. Thus if B is a minimum density set and
 215 $\frac{\tau(B)}{\nu(A) - \nu(A \cup B)} \leq 1$, then B satisfies the Density Condition; otherwise, no such B exists.

► **Theorem 5.** *The GREEDY ALGORITHM achieves approximation ratio*

$$1 + \frac{\tau^*}{\text{opt}} \ln \frac{\nu_0 - \nu^*}{\tau^*} = 1 + \frac{\tau^*}{\text{opt}} \cdot \ln \left(1 + \frac{\nu_0 - \text{opt}}{\tau^*} \right).$$

Proof. Let ℓ be the number of iterations. Let $A_0 = \emptyset$ and for $i = 1, \dots, \ell$ let A_i be the
 intermediate solution at the end of iteration i and $B_i = A_i \setminus A_{i-1}$. Let $\nu_i = \nu(A_i)$, $i = 0, \dots, \ell$.
 Then:

$$\frac{\tau(B_i)}{\nu_{i-1} - \nu_i} \leq \min \left\{ 1, \frac{\tau^*}{\nu_{i-1} - \nu^*} \right\} \quad i = 1, \dots, \ell$$

Since ν is decreasing

$$\sum_{i=1}^{\ell} \tau(B_i) \leq \sum_{i=1}^{\ell} \min \left\{ 1, \frac{\tau^*}{\nu_{i-1} - \nu^*} \right\} (\nu_{i-1} - \nu_i)$$

This is the lower Darboux sum of the function $f(\nu) = \begin{cases} 1 & \text{if } \nu \leq \tau^* + \nu^* \\ \frac{\tau^*}{\nu - \nu^*} & \text{if } \nu > \tau^* + \nu^* \end{cases}$ in the
 interval $[\nu_\ell, \nu_0]$ w.r.t. the partition $\nu_\ell < \nu_{\ell-1} < \dots < \nu_0$. We claim that $\tau^* + \nu^* \geq \nu_\ell$. For
 this, note that $\frac{\tau(A^*)}{\nu(A) - \nu(A \cup A^*)} \geq 1$, thus since ν is decreasing $\tau(A^*) \geq \nu(A) - \nu(A \cup A^*) \geq$
 $\nu(A) - \nu(A^*)$. Consequently, $\sum_{i=1}^{\ell} \tau(B_i)$ is bounded by

$$\int_{\nu_\ell}^{\nu_0} f(\nu) d\nu = \int_{\nu_\ell}^{\tau^* + \nu^*} 1 d\nu + \int_{\tau^* + \nu^*}^{\nu_0} \frac{\tau^*}{\nu - \nu^*} d\nu = \tau^* + \nu^* - \nu_\ell + \tau^* \ln \frac{\nu_0 - \nu^*}{\tau^*}$$

Let $A = \bigcup_{i=1}^{\ell} B_i$ be the set computed by the algorithm. Since τ is sub-additive

$$\tau(A) \leq \sum_{i=1}^{\ell} \tau(B_i) \leq \tau^* + \nu^* - \nu(A) + \tau^* \ln \frac{\nu_0 - \nu^*}{\tau^*}$$

216 Thus the approximation ratio is bounded by $\frac{\tau(A) + \nu(A)}{\text{opt}} \leq 1 + \frac{\tau^*}{\text{opt}} \ln \frac{\nu_0 - \nu^*}{\tau^*}$. ◀

3 Algorithm for general thresholds (Theorem 1)

Given an instance $G = (V, E)$, \mathbf{t} , R of ACTIVATION EDGE-COVER the corresponding GENERALIZED MIN-COVERING instance U, τ, ν is defined as follows. We put at each node $u \in V$ a large set of “assignment units”, and let U be the union of these sets of “assignment units”. Note that to every $A \subseteq U$ naturally corresponds the assignment \mathbf{a} where a_u is the number of units in A put at u . It would be more convenient to define ν and τ in terms of assignments, by considering instead of a set $A \subseteq U$ the corresponding assignment \mathbf{a} .

To define ν and τ , let us recall the assignments \mathbf{q} and \mathbf{c} from the Introduction. We have $c_v = q_v = 0$ if $v \in V \setminus R$ and for $u \in R$:

- $q_u = \min_{e=uv \in E} t_u^e$ is the minimum threshold at u of an edge in E incident to u .
- $c_u = \min_{e=uv \in E} (t_u^e + t_v^e) - q_u$, so $c_u + q_u$ is the minimum value of an edge in E incident to u .

We let $Q = \mathbf{q}(V) = \mathbf{q}(R)$ and $C = \mathbf{c}(R)$. Note that $\mathbf{c}(R') \leq \theta \mathbf{q}(R')$ for any $R' \subseteq R$; in particular, $C \leq \theta Q$. For an assignment \mathbf{a} that “augments” \mathbf{q} let $R_{\mathbf{q}+\mathbf{a}}$ denote the set of terminals covered by $E_{\mathbf{q}+\mathbf{a}}$. A natural definition of the potential and the payment functions would be $\tau(\mathbf{a}) = \mathbf{a}(V)$ and $\nu(\mathbf{a}) = (\mathbf{c} + \mathbf{q})(R \setminus R_{\mathbf{q}+\mathbf{a}})$ but this will enable to prove only ratio $1 + \ln(\theta + 1)$. We show a better ratio by adding to the potential in advance the “fixed” part Q . We define

$$\tau(\mathbf{a}) = \mathbf{a}(V) \quad \nu(\mathbf{a}) = Q + \mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}})$$

It is easy to see that ν is decreasing, τ is sub-additive, and $\tau(\mathbf{0}) = 0$.

The next lemma shows that the obtained GENERALIZED MIN-COVERING instance is equivalent to the original ACTIVATION EDGE-COVER instance.

► **Lemma 6.** *If $\mathbf{q} + \mathbf{a}$ is a feasible solution for ACTIVATION EDGE-COVER then $\tau(\mathbf{a}) + \nu(\mathbf{a}) = Q + \mathbf{a}(V)$. If \mathbf{a} is a feasible solution for GENERALIZED MIN-COVERING then one can construct in polynomial time a feasible solution for ACTIVATION EDGE-COVER of value at most $\tau(\mathbf{a}) + \nu(\mathbf{a})$. In particular, both problems have the same optimal value, and GENERALIZED MIN-COVERING has an optimal solution \mathbf{a}^* such that $\nu(\mathbf{a}^*) = Q$ and thus $\text{opt} = \tau(\mathbf{a}^*) + Q$.*

Proof. If $\mathbf{q} + \mathbf{a}$ is a feasible ACTIVATION EDGE-COVER solution then $R_{\mathbf{q}+\mathbf{a}} = R$ and thus $\nu(\mathbf{a}) = Q$. Consequently, $\tau(\mathbf{a}) + \nu(\mathbf{a}) = \mathbf{a}(V) + Q$.

Let now \mathbf{a} be a GENERALIZED MIN-COVERING solution. The assignment $\mathbf{q} + \mathbf{a}$ has value $Q + \mathbf{a}(V)$ and activates the edge set $E_{\mathbf{q}+\mathbf{a}}$ that covers $R_{\mathbf{q}+\mathbf{a}}$. To cover $R \setminus R_{\mathbf{q}+\mathbf{a}}$, pick for every $u \in R \setminus R_{\mathbf{q}+\mathbf{a}}$ an edge uv with $t_u^{uv} + t_v^{uv}$ minimum. Let \mathbf{b} be an assignment defined by $b_u = c_u$ if $u \in R \setminus R_{\mathbf{q}+\mathbf{a}}$ and $b_u = 0$ otherwise. The set of picked edges can be activated by an assignment $\mathbf{q} + \mathbf{b}$ that has value $Q + \mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}})$. The assignment $\mathbf{q} + \mathbf{a} + \mathbf{b}$ activates both edge sets and has value $Q + \mathbf{a}(V) + \mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}}) = \tau(\mathbf{a}) + \nu(\mathbf{a})$, as required. ◀

For the obtained GENERALIZED MIN-COVERING instance, let us fix an optimal solution \mathbf{a}^* as in Lemma 6, so $\nu^* = Q$ and $\text{opt} = \tau^* + Q$. Denote $\nu_0 = \nu(\mathbf{0}) = Q + \mathbf{c}(R)$, and note that $\mathbf{c}(R) \leq \theta Q$. To apply Theorem 5 we need several bounds given in the next lemma.

► **Lemma 7.** $\frac{\text{opt}}{\tau^*} \geq 1 + \frac{1}{\theta}$, $\frac{\nu_0}{\tau^*} \leq (\theta + 1) \left(\frac{\text{opt}}{\tau^*} - 1 \right)$, and $\frac{\nu_0 - \nu^*}{\tau^*} \leq \Delta + 1$.

Proof. Note that

$$\tau^* + Q = \text{opt} \leq \nu_0 \leq (\theta + 1)Q .$$

In particular, $Q \geq \tau^*/\theta$, and this implies the first bound of the lemma

$$\frac{\text{opt}}{\tau^*} = 1 + \frac{Q}{\tau^*} \geq 1 + \frac{1}{\theta} .$$

23:8 Approximating activation edge-cover and facility location problems

248 The second bound of the lemma holds since $\nu_0 \leq (\theta + 1)Q = (\theta + 1)(\text{opt} - \tau^*)$.

249 The last bound of the lemma is equivalent to the bound $\mathbf{c}(R) \leq \tau^*(\Delta + 1)$. Let J be an
 250 inclusion minimal edge cover of R activated by $\mathbf{q} + \mathbf{a}^*$. Then J is a collection \mathcal{S} of node
 251 disjoint rooted stars with leaves in R . Let $S \in \mathcal{S}$. By the definition of \mathbf{c} , $\mathbf{a}^*(S) \geq \max_{u \in R \cap S} c_u$,
 252 thus $\mathbf{c}(R \cap S) \leq |R \cap S| \mathbf{a}^*(S) \leq (\Delta + 1) \mathbf{a}^*(S)$. Consequently, $\mathbf{c}(R) = \sum_{S \in \mathcal{S}} \mathbf{c}(R \cap S) \leq$
 253 $(\Delta + 1) \sum_{S \in \mathcal{S}} \mathbf{a}^*(S) \leq (\Delta + 1) \mathbf{a}^*(V)$. ◀

We will show later that the GREEDY ALGORITHM can be implemented in polynomial time; now we focus on showing that it achieves the approximation ratios stated in Theorem 1. Substituting Lemma 7 second bound in Theorem 5 second bound and denoting $x = \frac{\text{opt}}{\tau^*} - 1$, we get that $x \geq 1/\theta$ and that the ratio is bounded by

$$1 + \frac{\tau^*}{\text{opt}} \cdot \ln \left(1 + \frac{\nu_0}{\tau^*} - \frac{\text{opt}}{\tau^*} \right) \leq 1 + \frac{\ln(\theta x)}{x + 1} = 1 + f(x)$$

Consequently, the the ratio is bounded by $1 + \max\{f(x) : x \geq 1/\theta\}$. We now derive a formula for the maximum. We have $\lim_{x \rightarrow \infty} f(x) = 0$ (this can be shown using L'Hospital's Rule), and $f(1/\theta) = 0$. Also:

$$f'(x) = -\frac{1}{(x + 1)^2} \ln(\theta x) + \frac{1}{x + 1} \frac{1}{x}$$

254 Hence $f'(x) = 0$ if and only if $\frac{1}{x+1} \ln(\theta x) = \frac{1}{x}$, namely, $x + 1 = x \ln(\theta x)$. For the analysis,
 255 we substitute $x \leftarrow 1/x$, and get the equation $1 + x = \ln(\theta/x)$, where $0 < x \leq \theta$. Since
 256 the function $x + 1$ is strictly increasing and the function $\ln(\theta/x)$ is strictly decreasing, this
 257 equation has at most one root; we claim that this root exists and is in the interval $(0, \theta]$. To
 258 see this consider the function $h(x) = x + 1 - \ln(\theta/x)$, and note that h is continuous and that
 259 $h(\theta) = \theta + 1 > 0$ while $h(\epsilon) = \epsilon + 1 - \ln(\theta/\epsilon) < 0$ for $\epsilon > 0$ small enough.

260 From this we get that the ratio is bounded by $1 + \omega(\theta)$, where $\omega(\theta)$ is the root of the
 261 equation $x + 1 = \ln(\theta/x)$.

262 Substituting Lemma 7 third bound in Theorem 5 first bound and observing that $\tau^* \leq \text{opt}$
 263 we get that the ratio is bounded by $1 + \ln(\Delta + 1)$. In the case when R is an independent set
 264 in G , it is easy to see that Lemma 7 third bound improves to $\frac{\nu_0 - \nu^*}{\tau^*} \leq \Delta$, and we get ratio
 265 $1 + \ln \Delta$ in this case.

266 Finally, we show that the GREEDY ALGORITHM algorithm can be implemented in
 267 polynomial time. As was mentioned in Section 2 before Theorem 5, we just need to perform
 268 in polynomial time the following two operations for any assignment \mathbf{a} : to check the condition
 269 $\nu(\mathbf{a}) > \nu^*$, and to find an augmenting assignment \mathbf{b} of minimum density.

270 It is easy to see that assignments \mathbf{q} and \mathbf{c} can be computed in polynomial time, and thus
 271 the potential $\nu(\mathbf{a}) = Q + \mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}})$ can be computed in polynomial time, for any \mathbf{a} . Let
 272 \mathbf{a}^* be an optimal solution as in Lemma 6, and denote $\tau^* = \tau(\mathbf{a}^*)$ and $\nu^* = \nu(\mathbf{a}^*) = Q$. Then
 273 the condition $\nu(A) > \nu^*$ is equivalent to $\nu(\mathbf{a}) \geq Q$ and thus can be checked in polynomial
 274 time.

Now we show how to find an augmenting assignment \mathbf{b} of minimum density. Note that the density of an assignment \mathbf{b} w.r.t. \mathbf{a} is

$$\frac{\tau(\mathbf{b})}{\nu(\mathbf{a}) - \nu(\mathbf{a} + \mathbf{b})} = \frac{\mathbf{b}(V)}{\mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}}) - \mathbf{c}(R \setminus R_{\mathbf{q}+\mathbf{a}+\mathbf{b}})} = \frac{\mathbf{b}(V)}{\mathbf{c}(R_{\mathbf{q}+\mathbf{a}+\mathbf{b}} \setminus R_{\mathbf{q}+\mathbf{a}})}.$$

275 **► Lemma 8.** *There exists a polynomial time algorithm that given an instance of ACTIVATION*
 276 *EDGE-COVER and an assignment \mathbf{a} finds an assignment \mathbf{b} of minimum density.*

277 **Proof.** A **star** is a rooted tree $S = (V_S, E_S)$ with at least one edge such that only its root
 278 s may have degree ≥ 2 . We say that a star S is a **proper star** if all the leaves of S are
 279 terminals. We denote the terminals in S by $R_S = R \cap V_S$.

Since \mathbf{q}, \mathbf{a} are given assignments, we may simplify the notation by assuming that $R \leftarrow R \setminus R_{\mathbf{q}+\mathbf{a}}$ is our set of terminals, and that $\mathbf{a} \leftarrow \mathbf{q} + \mathbf{a}$ is our given assignment. Then the density of \mathbf{b} is just $\frac{\mathbf{b}(V)}{\mathbf{c}(R_{\mathbf{a}+\mathbf{b}})}$. Let \mathbf{b}^* be an assignment of minimum density, and let $J^* \subseteq E_{\mathbf{a}+\mathbf{b}^*}$ be an inclusion minimal $R_{\mathbf{a}+\mathbf{b}^*}$ -cover. Then J^* decomposes into a collection \mathcal{S} of node disjoint proper stars that collectively cover $R_{\mathbf{a}+\mathbf{b}^*}$. For $S \in \mathcal{S}$ let \mathbf{b}^S be the optimal assignment such that $\mathbf{a} + \mathbf{b}^S$ activates S . Since the stars in \mathcal{S} are node disjoint

$$\sum_{S \in \mathcal{S}} \mathbf{b}^S(V) \leq \mathbf{b}^*(V) \quad \text{and} \quad \sum_{S \in \mathcal{S}} \mathbf{c}(R_S) = \mathbf{c}(R_{\mathbf{a}+\mathbf{b}^*}).$$

280 By an averaging argument, $\frac{\mathbf{b}^S(V)}{\mathbf{c}(R_S)} \leq \frac{\mathbf{b}^*(V)}{\mathbf{c}(R_{\mathbf{a}+\mathbf{b}^*})}$ holds for some $S \in \mathcal{S}$, and since \mathbf{b}^* is a
 281 minimum density assignment, so is \mathbf{b}^S , and $\frac{\mathbf{b}^S(V)}{\mathbf{c}(R_S)} = \frac{\mathbf{b}^*(V)}{\mathbf{c}(R_{\mathbf{a}+\mathbf{b}^*})}$ holds. Consequently, it is
 282 sufficient to show how to find in polynomial time an assignment \mathbf{b} such that $\mathbf{a} + \mathbf{b}$ activates
 283 a proper star S and $\frac{\mathbf{b}(V)}{\mathbf{c}(R_S)}$ is minimal.

284 We may assume that we know the root v and the value $w = b_v$ of an optimal density pair
 285 S, \mathbf{b} ; there are at most $|V||E|$ choices and we can try all and return the best outcome. Let
 286 $R_w = \{u \in R : \text{there is a } uv\text{-edge } e \text{ with } t_v^e \leq a_v + w\}$. For $u \in R_w$ let b_u be the minimal
 287 non-negative number for which there is a uv -edge e with $a_v + w \geq t_v^e$ and $a_u + b_u \geq t_u^e$.
 288 Then our problem is equivalent to finding $R_S \subseteq R_w$ with $\sigma(R_S) = \frac{w + \mathbf{b}(R_S)}{\mathbf{c}(R_S)}$ minimum. This
 289 problem can be solved in polynomial time, by starting with $R_S = \emptyset$ and while there is
 290 $u \in R_w \setminus R_S$ with $\sigma(R_S + u) < \sigma(R_S)$, adding $u \in R_w \setminus R_S$ to R_S with b_u/c_u minimum. ◀


291 The proof of Theorem 1 is complete.

292 4 Locally uniform thresholds (Theorem 3)

293 Here we consider the BIPARTITE ACTIVATION EDGE-COVER problem with locally uniform
 294 thresholds. This means that each non-terminal $v \in V \setminus R$ has weight w_v and all edges incident
 295 to v have the same threshold t^v ; in the θ -bounded version $w_v \leq \theta t^v$. We consider a natural
 296 greedy algorithm that repeatedly picks a star S that minimizes the average price paid for
 297 each terminal (the quotient of the optimal activation value of S over $|R_S|$), and then removes
 298 R_S . Each time we choose a star S we distribute its activation value uniformly among its
 299 terminals, paying in the computed solution the average price for each terminal of S .

We now apply a standard “set-cover” analysis, cf. [24]. In some optimal solution fix an inclusion maximal star S^* with center v and terminals R_{S^*} covered by the algorithm in the order r_k, r_{k-1}, \dots, r_1 , where r_k is covered first and r_1 last; we bound the algorithm payment for covering R_{S^*} . Note that $1 \leq k \leq \Delta$. Denote $w = w_v$ and let t be the threshold of the terminals in S^* . Let S_i^* be the substar of S^* with leaves r_i, \dots, r_1 . At the start of the iteration in which the algorithm covers r_i , the terminals of S_i^* are uncovered. Thus the algorithm pays for covering r_i at most the average price paid by S_i^* , namely $(w + it)/i = w/i + t$. Over all iterations, the algorithm pays for covering R_{S^*} at most $wH_k + kt$, while the optimum pays $w + kt$. Thus the quotient between them is bounded by

$$\frac{wH_k + kt}{w + kt} = \frac{w/tH_k + k}{w/t + k} \leq \frac{\theta H_k + k}{\theta + k} = 1 + \frac{\theta(H_k - 1)}{\theta + k} \leq 1 + \max_{1 \leq k \leq \Delta} \frac{H_k - 1}{1 + k/\theta}.$$



■ **Figure 1** Tight example of ratio $\frac{73}{60}$ for unit thresholds.

Since any optimal solution decomposes into node disjoint stars, the last term bounds the approximation ratio, concluding the proof of Theorem 3. We make some observations about this bound. Let $g(k) = \frac{\theta(H_k - 1)}{\theta + k}$. We have

$$g(k + 1) - g(k) = \frac{\theta}{(k + \theta)(k + 1 + \theta)} \left(2 - H_k + \frac{\theta - 1}{k + 1} \right)$$

Thus $g(k + 1) \geq g(k)$ if and only if $2 - H_k + \frac{\theta - 1}{k + 1} > 0$. Hence if k_θ is the smallest integer such that $H_k \geq 2 + \frac{\theta - 1}{k + 1}$ then $\max_{1 \leq k \leq \Delta} g(k) = g(\min\{k_\theta, \Delta\})$. We do not have a more convenient formula of $\max_{k \geq 1} g(k)$ for arbitrary θ , but we can bound it using the inequality $H_k - 1 \leq \ln k$. Then we have:

$$\max_{1 \leq k \leq \Delta} \frac{\theta(H_k - 1)}{k + \theta} \leq \max_{x \geq 1} \frac{\theta \ln x}{x + \theta} = \max_{x \geq 1} f(x).$$

Using fundamental calculus one can see that the maximum is attained when $x + \theta = x \ln x$, and substituting this in $f(x)$ we get that $\max_{x \geq 1} f(x) = \frac{\theta}{\alpha}$ where α is the solution to the equation $x + \theta = x \ln x$. We have $\alpha = \frac{\theta}{W(\theta/e)} = \frac{\theta}{\omega(\theta)}$. Thus the ratio is bounded by $1 + \omega(\theta)$, a bound that we got before in Theorem 1.

If $\theta = 1$ then $k_\theta = 4$, since $H_3 = 11/6 < 2$ and $H_4 = 25/12 > 2$. We have $g(4) = \frac{13}{60}$, so for $\theta = 1$ we get ratio $1 + g(4) = \frac{73}{60}$. The ratio $\frac{73}{60}$ is tight for unit thresholds, as shows the example in Fig. 1. The instance has 48 terminals (in black), and two sets of covering nodes: the upper 12 nodes that form an optimal cover, and the bottom 13 nodes. The bottom nodes have 3 nodes of degree 4, 4 of degree 3, and 6 of degree 2. The algorithm may start taking all bottom nodes, and only then add the upper ones, thus creating a solution of value 73, instead of the optimum 60.

5 Unit thresholds (Theorem 4)

Here we consider the case of unit thresholds when $t_u^e = t_v^e = 1$ for every uv -edge e . By a reduction from [3], we may assume that the instance is bipartite. Specifically, for any optimal assignment \mathbf{a} we have $a_u = 1$ for all $u \in R$, hence we can consider the residual instance obtained by removing the terminals covered by edges with both ends in R ; in the new obtained instance R is an independent set, and recall that we may assume that $V \setminus R$ is an independent set.

One can observe that in the obtained bipartite instance, \mathbf{a} is an optimal solution if and only if $a_v \in \{0, 1\}$ for all $v \in V$, $a_v = 1$ for all $v \in R$, and the set $C = \{v \in V \setminus R : a_v = 1\}$ covers R , meaning that R is the set of neighbors of C . Namely, our problem is equivalent to $\min\{|C| + |R| : C \subseteq V \setminus R, C \text{ covers } R\}$. On the other hand the problem $\min\{|C| : C \subseteq V \setminus R, C \text{ covers } R\}$ is essentially the (unweighted) SET-COVER problem, and C is a feasible solution to this SET-COVER instance if and only if $C \cup R$ is the characteristic set of a feasible assignment for the ACTIVATION EDGE-COVER instance. Note that both problems are equivalent w.r.t. their optimal solutions but may differ w.r.t. approximation ratios, since if C^* is an optimal SET-COVER solution then $\frac{|C| + |R|}{|C^*| + |R|}$ may be much smaller than $\frac{|C|}{|C^*|}$.

327 Recall that a standard greedy algorithm for SET-COVER repeatedly picks the center
 328 of a largest star and removes the star from the graph. This algorithm has ratio H_k for
 329 k -SET-COVER, where $k = \Delta$ is the maximum degree of a non-terminal (the maximum size
 330 of a set). However, the same algorithm achieves a much smaller ratio $\frac{73}{60}$ for ACTIVATION
 331 EDGE-COVER with unit thresholds; the ratio $\frac{73}{60}$ was established in [3], and it also follows
 332 from the case $\theta = 1$ in Theorem 3. In what follows we denote by α_k the best known ratio for
 333 k -SET-COVER. We have $\alpha_1 = \alpha_2 = 1$ ($k = 2$ is the EDGE-COVER problem) and $\alpha_3 = 4/3$ [6].
 334 The current best ratios for $k \geq 4$ are due to [9] (see also [14, 1]). We summarize the current
 335 values of α_k for $k \leq 7$ in the following table.

α_1	α_2	α_3	α_4	α_5	α_6	α_7
1	1	$\frac{4}{3}$	$\frac{73}{48}$	$\frac{26}{15}$	$\frac{28}{15}$	$\frac{212}{105}$

■ **Table 2** Current values of α_k for $k \leq 7$.

336 We now show how these ratios for k -SET-COVER can be used to approximate the
 337 ACTIVATION EDGE-COVER problem with unit costs. We start by describing a simple
 338 algorithm with ratio $1 \frac{67}{360} < \frac{73}{60}$, that uses only the $k = 2$ case.

339 **Algorithm 1:** ratio $1 \frac{67}{360}$

```

340 1  $A \leftarrow \emptyset$ 
341 1 while there exists a star with at least 3 terminals do
342 2   add to  $A$  and remove from  $G$  the node-set of a maximum size star
343 3   add to  $A$  an optimal solution of the residual instance

```

We claim that the above algorithm achieves approximation ratio $1 \frac{67}{360}$ for ACTIVATION EDGE-COVER (a similar analysis implies ratio $H_k - \frac{1}{6}$ for SET-COVER). In some optimal solution fix a star S^* with terminals covered in the order r_k, r_{k-1}, \dots, r_1 , where r_k is covered first and r_1 last; we bound the algorithm payment to cover these terminals. Let S_i^* be the substar of S^* with leaves r_i, \dots, r_1 . At the start of the iteration when r_i is covered, the terminals of S_i^* are uncovered. Thus the algorithm pays for covering r_i at most the density of S_i^* , namely, $(i+1)i = 1 + 1/i$. Over all iterations, the algorithm pays for covering R_S at most $k + H_k$, while the optimum pays $k + 1$. If $k = 1$ then the algorithm pays at most the amount of the optimum. We claim that if $k \geq 2$ then in fact the payment is at most $k + H_k - 1/6$. If $k = 2$ then the payment is at most $3 < 3 + H_3 - 1/6$ (we pay 3 if the star “survives” all the iterations before the last). For $k \geq 3$, the pay for the last 3 terminals is either: $4/3$ for each of for r_3, r_2 and 2 for r_1 (a total of $14/3$), or $4/3$ for r_3 and 3 for r_2, r_1 (a total of $13/3$). The maximum is $14/3 = 3 + H_3 - 1/6$. Consequently, the ratio is bounded by

$$\max_{k \geq 2} \frac{k + H_k - 1/6}{k + 1} = 1 + \max_{k \geq 2} \frac{H_k - 7/6}{k + 1} = 1 + \max_{k \geq 2} g(k)$$

344 By fundamental computations we have $g(k+1) - g(k) = \frac{13/6 - H_k}{(k+1)(k+2)}$. Thus $g(k)$ is increasing
 345 iff $H_k < \frac{13}{6}$. Since $H_4 = \frac{23}{12} < \frac{13}{6}$ and $H_5 = \frac{137}{60} > \frac{13}{6}$, we get that $\max_{k \geq 2} g(k) = g(5) = \frac{67}{360}$,
 346 so we have ratio $1 \frac{67}{360}$.

347
 348 We now show ratio $\frac{8}{7} < \frac{1555}{1347} < \frac{7}{6}$. As in the greedy algorithm for SET-COVER, we
 349 repeatedly remove an inclusion maximal set of disjoint stars with maximum number of leaves
 350 and pick the set of roots of these stars. The difference is that each time stars with more

23:12 Approximating activation edge-cover and facility location problems

351 than k leaves are exhausted, we compute an α_k -approximate solution A_k for the remaining
 352 k -SET-COVER instance; we let $A_0 = \emptyset$. This gives many SET-COVER solutions, each is a
 353 union of the centers of stars picked and A_k ; we choose the smallest one, and together with R
 354 this gives a feasible ACTIVATION EDGE-COVER solution. Formally, the algorithm is:

355 **Algorithm 2:** ratio $\rho = \frac{1555}{1347} < 1.1545$

356 **1** for $k \leftarrow \Delta$ downto 0 do
 357 **2** remove from G a maximal collection of node disjoint $(k+1)$ -stars
 358 let C_{k+1} be the set of the roots of the stars removed so far
 359 **3** compute an α_k -approximate k -SET-COVER solution A_k in G
 360 **4** return the smallest set $C_{k+1} \cup A_k$, $k \in \{\Delta, \dots, 0\}$

361 Since we claim ratio $\frac{1555}{1347} < \frac{8}{7}$, at iterations when $k \geq 7$ step 3 can be skipped, since then
 362 we can apply a standard “local ratio” analysis [2]. Indeed, when a star with $k \geq 7$ terminals
 363 is removed, the partial solution value increases by $k+1$ while the optimum decreases by at
 364 least k . Hence for $k \geq 7$ it is a $\frac{k+1}{k} \leq \frac{8}{7}$ local ratio step. Consequently, we may assume that
 365 $\Delta \leq 6$, provided that we do not claim ratio better than $8/7$.

366 Let $r = |R|$. Let τ be the optimal value to the initial SET-COVER instance. At iteration
 367 k the algorithm computes a solution of value at most $\alpha_k \tau + r + |C_{k+1}|$. Thus we get ratio ρ
 368 if $\rho(r + \tau) \geq \alpha_k \tau + r + |C_{k+1}|$ holds for some $k \leq 6$. Otherwise,

$$\begin{aligned} 369 \quad \rho(r + \tau) &< \alpha_6 \tau + r \\ 370 \quad \rho(r + \tau) &< \alpha_5 \tau + r + |C_6| \\ 371 \quad \rho(r + \tau) &< \alpha_4 \tau + r + |C_5| \\ 372 \quad \rho(r + \tau) &< \alpha_3 \tau + r + |C_4| \\ 373 \quad \rho(r + \tau) &< \alpha_2 \tau + r + |C_3| \\ 374 \quad \rho(r + \tau) &< \alpha_1 \tau + r + |C_2| \\ 375 \quad \rho(r + \tau) &< r + |C_1| \end{aligned}$$

376 Denote $\sigma = \alpha_1 + \dots + \alpha_5 = \frac{1581}{240}$. Note that $|C_1| + \dots + |C_6| = r$, since in this sum the
 377 number of stars with k leaves is summed exactly k times, $k = 1, \dots, 6$. The first inequality,
 378 and the inequality obtained as the sum of the other six inequalities gives the following two
 379 inequalities:

$$\begin{aligned} 380 \quad \rho(r + \tau) &< \alpha_6 \tau + r \\ 381 \quad 6\rho(r + \tau) &< \sigma \tau + 7r \end{aligned}$$

382 Dividing both inequalities by τ and denoting $x = r/\tau$ gives:

$$\begin{aligned} 383 \quad \rho(x + 1) &< \alpha_6 + x \\ 384 \quad 6\rho(x + 1) &< \sigma + 7x \end{aligned}$$

Since $\rho > 1$ and $7 > 6\rho$ this is equivalent to:

$$\frac{6\rho - \sigma}{7 - 6\rho} < x < \frac{\alpha_6 - \rho}{\rho - 1}$$

We obtain a contradiction if ρ is the solution of the equation $\frac{6\rho - \sigma}{7 - 6\rho} = \frac{\alpha_6 - \rho}{\rho - 1}$, namely

$$\rho = \frac{7\alpha_6 - \sigma}{6\alpha_6 - \sigma + 1} = 1 + \frac{\alpha_6 - 1}{6\alpha_6 - \sigma + 1} = 1 + \frac{208}{1347} = \frac{1555}{1347}.$$

385 This concludes the proof of Theorem 4.

386 — **References** —

- 387 1 S. Athanassopoulos, I. Caragiannis, and C. Kaklamanis. Analysis of approximation algorithms
388 for k -set cover using factor-revealing linear programs. *Theory Comput. Syst.*, 45(3):555–576,
389 2009.
- 390 2 Reuven Bar-Yehuda. One for the price of two: A unified approach for approximating covering
391 problems. *Algorithmica*, 27(2):131–144, 2000.
- 392 3 G. Calinescu, G. Kortsarz, and Z. Nutov. Improved approximation algorithms for minimum
393 power covering problems. In *WAOA*, pages 134–148, 2018.
- 394 4 K. Chandrasekaran, R. M. Karp, E. Moreno-Centeno, and S. Vempala. Algorithms for implicit
395 hitting set problems. In *SODA*, pages 614–629, 2011.
- 396 5 V. Chvatal. Greedy heuristic for the set-covering problem. *Mathematics of Operations Research*,
397 4(3):233–235, 1979.
- 398 6 Rong-chii Duh and Martin Fürer. Approximation of k -set cover by semi-local optimization. In
399 *STOC*, pages 256–264, 1997.
- 400 7 J. Edmonds. Paths, trees, and flowers. *Canadian Journal of Mathematics*, 17:449–467, 1965.
- 401 8 U. Feige. A threshold of $\ln n$ for approximating set cover. *J. of the ACM*, 45(4):634–652, 1998.
- 402 9 M. Fürer and H. Yu. Packing-based approximation algorithm for the k -set cover problem. In
403 *ISAAC*, pages 484–493, 2011.
- 404 10 C. Gröpl, S. Hougardy, T. Nierhoff, and H. J. Prömel. Approximation algorithms for the
405 steiner tree problem in graphs. In X. Z. Cheng and D-. Z. Du, editors, *Steiner Trees in*
406 *Industry*, pages 235–279. Kluwer Academic Publishers, 2001.
- 407 11 M. Hajiaghayi, G. Kortsarz, V. Mirrokni, and Z. Nutov. Power optimization for connectivity
408 problems. *Math. Programming*, 110(1):195–208, 2007.
- 409 12 D. S. Johnson. Approximation algorithms for combinatorial problems. *J. Comput. Syst. Sci.*,
410 9(3):256–278, 1974.
- 411 13 G. Kortsarz and Z. Nutov. Approximating minimum-power edge-covers and 2, 3-connectivity.
412 *Discrete Applied Mathematics*, 157:1840–1847, 2009.
- 413 14 A. Levin. Approximating the unweighted k -set cover problem: greedy meets local search. In
414 *WAOA*, pages 290–301, 2006.
- 415 15 Z. Nutov. Activation network design problems. In T. F. Gonzalez, editor, *Handbook on*
416 *Approximation Algorithms and Metaheuristics, Second Edition*, volume 2, chapter 15. Chapman
417 & Hall/CRC, 2018.
- 418 16 D. Panigrahi. Survivable network design problems in wireless networks. In *SODA*, pages
419 1014–1027, 2011.
- 420 17 R. Raz and S. Safra. A sub constant error probability low degree test, and a sub constant
421 error probability PCP characterization of NP. In *STOC*, pages 475–484, 1997.
- 422 18 G. Robins and A. Zelikovsky. Tighter bounds for graph steiner tree approximation. *SIAM J.*
423 *Discrete Math.*, 19(1):122–134, 2005.
- 424 19 A. Schrijver. *Combinatorial Optimization, Polyhedra and Efficiency*. Springer-Verlag Berlin,
425 Heidelberg New York, 2004.
- 426 20 D B. Shmoys. Approximation algorithms for facility location problems. In *APPROX*, pages
427 265–274, 2000.
- 428 21 P. Slavik. A tight analysis of the greedy algorithm for set cover. *J. Algorithms*, 25(2):237–254,
429 1997.
- 430 22 J. Vygen. Approximation algorithms for facility location problems (lecture notes). Technical
431 Report 05950, Research Institute for Discrete Mathematics, University of Bonn, 2005.
- 432 23 L. Wolsey. An analysis of the greedy algorithm for the submodular set covering problem.
433 *Combinatorica*, 2(4):385–393, 1982.
- 434 24 N. E. Young. Greedy set-cover algorithms. In *Encyclopedia of Algorithms*, pages 886–889.
435 Springer, 2016.